Page 1 of 9

#### SECTION A **MULTIPLE CHOICE Ouestion** 1

- In a galvanic cell, oxidation occurs at the
- A. anode which is the negative electrode.
- B. anode which is the positive electrode.
- C. cathode which is the negative electrode.
- D. cathode which is the positive electrode.

## **Question 2**

The correct order of weakest to strongest oxidant is


- A.  $\text{Sn}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Br}_2$ ,  $\text{Cr}_2\text{O}_7^{2-}$ B.  $\text{Fe}^{2+}$ ,  $\text{Sn}^{2+}$ ,  $\text{Br}_2$ ,  $\text{Cr}_2\text{O}_7^{-2-}$

## **Ouestion 3**

The salt bridge of a galvanic cell balances the charges in the half cells by supplying

- A. cations to the cathode and anions to the anode.
- B. anions to the cathode and cations to the anode.
- electrons to the cathode and receiving electrons from the anode. C.
- D. electrons to the anode and receiving electrons from the cathode.

## **Question 4**



The diagram above shows the basic structure of a mercury-zinc button cell. The cell can be classified as an alkaline cell. The reaction occurring at the cathode could be written as

- A.  $Zn(s) + 2OH^{-}(aq) \longrightarrow ZnO(s) + H_{2}O(1) + 2e$
- B.  $ZnO(s) + H_2O(1) + 2e^- \longrightarrow Zn(s) + 2OH^-(aq)$
- C.  $Hg(1) + 2OH^{-}(aq) \longrightarrow HgO(s) + H_2O(1) + 2e$
- D.  $HgO(s) + H_2O(1) + 2e^- \longrightarrow Hg(1) + 2OH^-(aq)$

## **Ouestion 5**

During discharge of a lead-acid battery, the following occurs.

 $Pb(s) + PbO_2(s) + 2H_2SO_4(aq) \longrightarrow 2PbSO_4(s) + 2H_2O(1)$ 

The oxidant and the electrolyte are respectively

- A. lead(II) sulfate and water.
- B. lead and sulfuric acid.
- C. lead(IV) oxide and water..
- D. lead(IV) oxide and sulfuric acid.

## **Question 6**

A metal object to be electroplated is connected to the negative terminal of the power supply and suspended in an electrolyte containing copper(II) ions, silver(I) ions, gold(I) ions and magnesium(II) ions. The other inert electrode is connected to the positive terminal. The metal that initially deposits is

- A. copper.
- B. silver.
- C. gold.
- D. magnesium.

## **Ouestion 7**

Which of the following is not true of all operating fuel cells?

- A. Reactants are supplied continuously.
- B. The electrode must be porous and lined with catalysts.
- C. The products are pollutants and disposal systems have to be incorporated into the fuel cell design.
- D. They cannot be switched on and off like galvanic cells

## **Question 8**

As opposed to the galvanic cell, the electrolytic cell involves a

- A. non-spontaneous reaction and energy is required.
- B. spontaneous reaction and energy is required.
- C. non-spontaneous reaction and energy is released.
- D. spontaneous reaction and energy is released.

#### **Question 9**

Hydrogen bonding is not a significant factor in

- A. determining the primary structure of a protein molecule.
- B. accounting for the high solubility of glucose in water.
- C. determining the  $\alpha$ -helical shape of a protein molecule.
- D. determining the specificity of enzymes.

## **Question 10**

When the disaccharide sucrose reacts to form glucose and fructose, it undergoes a

- A. condensation reaction involving the breakage of an ester linkage.
- B. hydrolysis reaction involving the formation of an ether linkage.
- C. hydrolysis reaction with all products having a formula of  $C_6H_{12}O_6$ .
- D. reaction involving the formation of an ether linkage.

## **Question 11**

Amylase acts as an enzyme in the hydrolysis of the carbohydrate starch by

- A. decreasing the enthalpy of the reaction.
- B. decreasing the activation energy of the reaction.
- C. increasing the activation energy of the reaction.
- D. causing a different type of biochemical reaction.

## **Question 12**

The protein molecules in food are digested to form amino acids. Which one of the following statements about  $\alpha$ -amino acids is correct?

- A. Being molecules, they have low melting points.
- B. All amino acids have an amino functional group, -NH<sub>3</sub>.
- C. They are amphiprotic.
- D. They undergo hydrolysis to form polypeptides.

#### **Question 13**

When a glycine solution of pH 6.0 is placed in an electrolytic cell the glycine molecules do not migrate to either electrode. At this pH glycine mostly exists as

- A. NH<sub>3</sub><sup>+</sup>CH<sub>2</sub>COO<sup>+</sup>
- B. NH<sub>2</sub>CH<sub>2</sub>COOH
- C.  $NH_3^+CH_2COOH$
- D. Equal amounts of NH<sub>2</sub>CH<sub>2</sub>COO<sup>-</sup> and NH<sub>3</sub><sup>+</sup>CH<sub>2</sub>COOH

#### Question 14

Acid rain is partly due to the presence of  $SO_2$  in the atmosphere. One source of  $SO_2$  is the gases produced during combustion of fossil fuels. Fossil fuels are formed from the decomposition of plants. The molecules in the original plants that produced the  $SO_2$  are

- A. proteins
- B. carbohydrates
- C. unsaturated fats
- D. saturated fats.

#### **Question 15**

One step in radioactive decay of sodium is

- $^{22}_{11}Na \longrightarrow ^{22}_{10}Ne + X$
- Particle X is
- A. a proton
- B. a neutron
- C. electron
- D. a positron

#### Page 3 of 9

## Question 16

In our Sun,

- A. nuclear fission occurs, with the absorption of energy.
- B. nuclear fusion occurs, with the release of energy.
- C. nuclear fusion occurs, with the absorption of energy.
- D. nuclear fission occurs, with the release of energy.

## **Question 17**

When our Sun first formed, it's core became extremely hot because

- A. hydrogen combustion occurred there.
- B. electrons were stripped off hydrogen atoms.
- C. the hydrogen gas expanded.
- D. as gas particles were pulled into the core their kinetic energy increased.

## **Question 18**

In the original version of the Periodic Table devised by the Russian Dmitri Mendeleev, elements were arranged in order of increasing

- A. atomic number with elements of similar properties placed in vertical groups.
- B. atomic number with elements of similar properties placed in horizontal periods.
- C. atomic mass with elements of similar properties placed in vertical groups.
- D. atomic mass with elements of similar properties placed in horizontal periods.

## **Question 19**

The element with electronic configuration  $1s^22s^22p^63s^23p^63d^4s^2$  in the Periodic Table is

- A. in Group 2, Period 3, p block.
- B. in Group 3, Period 4, d block.
- C. in Group 2, Period 4, s block.
- D. in Group 4, Period 3, s block.

## **Question 20**

For the first five elements of the first-row transition metals, the maximum oxidation number occurs when all 4s and 3d electrons are lost. The maximum oxidation number of chromium is

- A. +3.
- B. +4.
- C. +5.
- D. +6.


# Question 1 (15 minutes, 13 marks)

#### Part A

In a commercial process called the Dow process, molten magnesium chloride is electrolysed and molten magnesium metal is produced.

- (a) On the following schematic diagram, label the boxes
  - anode and the cathode.
  - the polarity of the electrodes.
  - the direction of electron flow.

(3 marks)



- (b) Write the equation for the reaction that occurs at (i) the anode
  - (ii) the cathode

(c) Explain why

(i) molten and not aqueous magnesium chloride is used.

(ii) the products must be kept separate.

(2 marks)

(2 marks)

#### Part B

(a) Thomas Edison needed to find how much electricity his customers had used. One of his first solutions to this problem was to take a small fraction of that current and use it, to electrolyse a zinc salt solution and then to weigh the mass of zinc deposited. The cell was called a zinc `coulometer,. Calculate the mass of zinc that would be deposited in the month of November (30 days) if 2.0 mA of current passed through the coulometer.


#### Page 5 of 9

(b) A student electrolyses the following solutions using inert graphite electrodes. Predict the products likely to form at each electrode and write the formulae in the space provided.

(i)	1 M silver nitrate	Anode	Cathode
(ii)	1 M sodium iodide	Anode	Cathode
(iii)	1 M copper(II) bromide	Anode	Cathode

#### Question 2 (8 minutes, 7 marks)

(a) A flowchart of a nuclear power station is shown below. Below the flowchart describe the energy transformations that have occurred at Stages I, II and III.



#### Question 3 (8 minutes, 8 marks)

A health food company investigates a new product, 'Protofat'. This product was hoped to act as a synthetic food in third world countries. 'Protofat' is a combination of fat and protein exclusively. The energy content of this new product was determined by means of bomb calorimetry. A 0.298 g sample of 'Protofat' was placed in a bomb calorimeter and then allowed to react with excess oxygen gas. The results of the experiment were recorded.

Initial temperature of the bomb calorimeter and sample =  $17.234 \text{ }^{\circ}\text{C}$ 

Temperature after complete combustion of the sample = 20.514 °C

(3 marks)

(3 marks)

Pag	ge 6 of 9				
Ten	mperature after a 5.10 A current at 5.00 V = $21.202 \circ C$	2			
was	s passed through the heating coil for 1.02 minutes.				
(a)	Calculate the calibration factor for the calorimeter.				
				(2	marks)
(b)	Calculate the heat energy released by the sample.				
				(	1 mark)
(c)	What is the energy content of $Protofat'$ in kJ/g ?				
(d)	What is a dietary problem with using Protofat' as the only	ly food sourc	e in a person's di	iet?	(1 mark)
(e)	Given that protein has an energy content of 17 kJ/g and fat has the Protofat' sample.	an energy co	ntent of 37 kJ/g,	calculate the per-	(1 mark) centage of fat in
				(3 marks)	
The	estion 4 (8 minutes, 8 marks) e structural formula below is a segment of a molecule with a rela 23 000. The molecule was isolated from E. coli bacteria.	tive molecula	ar mass		
	H CH2OH H CH2OH         -CO-C-NH-C-NH		CH₂OH   CO-C-NH-CO	CH3   - C-NH-	
		 H	 H	 H	
(a)	On the diagram circle one linkage, which connects the monome	er units toget	her.		/1
(b) <sup>•</sup>	Write the name of this linkage.				(1 mark)

(1 mark)

#### Page 7 of 9

(c)	On the structural formula, circle one hydrogen atom that is capable of forming bonds associated with the secondary structure
	of the protein.

(1 mark)

(d) How many different amino acids are present in this section of the molecule

(e) What could be a possible biological function of this molecule?

(f) Name the molecules that are formed as the waste products in the complete digestion of this polypeptide molecule.

(1 mark) (g) The second amino acid in the polpeptide chain is serine residue. In the space below draw a structural formula of serine in a pH 3 solution.

Question 5 (5 minutes, 5 marks)

(a) Write the molecular formula for the reductant that is consumed in cellular respiration.

(b) Draw the structure of the bond linking fatty acids to glycerol.

(c) Give the formula of two products of N-fixing bacteria.

(d) Write the symbols for the elements contained in all carbohydrates.

(e) Draw the structural formula of glycerol.

(5 marks)

(2 marks)

#### Question 6 (12 minutes, 9 marks)

A British scientist Humphry Davy investigated the properties of many metals. In 1812, he wrote "Potassium and platinum, if we except their lustre, colour and power of conducting electricity, are bodies extremely dissimilar". (a) Davy noticed that potassium, a group I metal, is much less dense than platinum, a transitionmetal. Explain this difference.

(1 mark)

(1 mark)

(2 marks)

# Page 8 of 9

- (b) Platinum, in a similar reaction to gold, can be dissolved by solutions containing cyanide ions, CN<sup>-</sup>, to form the complex ion tetracyanoplatinate Pt(CN)<sub>4</sub><sup>2-</sup>.
  (i) What is the oxidation number of Pt in Pt(CN)<sub>4</sub><sup>2-</sup>

	(ii) <u> </u>	Praw a diagram showing the	e structure of $Pt(CN)_4^2$	•		_
					(.	3 marks)
c)		her metal studied by Davy w Vrite the electron configurat		he ground state.		
	(ii) V	Vrite the electronic configur	ration of oxygen with o	one electron in an ex	cited state.	
	(iii)	One compound containi manganese in this com		anese and oxygen is	$MnO_4^-$ . What is	the oxidation number of
	(iv)	Why is manganese able	to form compounds in	which it has differe	ent oxidation stat	les ?
	(iv)	Why is manganese able	to form compounds in	which it has differe	ent oxidation stat	(4 marks)
od	estion	7 (5 minutes, 4 marks) luminium and sulfur are thr two trends in the chemical	ree elements found in p	period 3.		
od	<b>estion</b> lium, a Name	7 (5 minutes, 4 marks) luminium and sulfur are thr two trends in the chemical	ree elements found in p	period 3.		(4 marks)
od	<b>estion</b> lium, a Name	7 (5 minutes, 4 marks) luminium and sulfur are thr two trends in the chemical	ree elements found in p	period 3.		(4 marks)
od	<b>estion</b> lium, a Name	7 (5 minutes, 4 marks) luminium and sulfur are thr two trends in the chemical	ree elements found in p	period 3.		(4 marks)
od	<b>estion</b> lium, a Name	7 (5 minutes, 4 marks) luminium and sulfur are thr two trends in the chemical	ree elements found in p	period 3.		(4 marks)
od a)	estion lium, a trends	7 (5 minutes, 4 marks) luminium and sulfur are thr two trends in the chemical s.	ree elements found in p properties of elements	period 3. s that can be seen ac	ross period 3 and	(4 marks)

(2 marks)

# Question 8 (5 minutes, 3 marks)

Hydrogen is the most abundant element in the Universe, over 90% by mass.

(a) Why is the mass of hydrogen in the universe decreasing?

		(1 mark)
(b)	How do we know the composition of elements in stars and other parts of the universe?	
(c)	Why can't our Sun produce the range of elements found on Earth? (1 mark)	

(1 mark)

END OF TASK.