



THE SCHOOL FOR EXCELLENCE
UNIT 4 SPECIALIST MATHEMATICS 2006
COMPLIMENTARY WRITTEN EXAMINATION 1 - SOLUTIONS

QUESTION 1

a. $\vec{CD} = \vec{CO} + \vec{OD} = -\vec{OC} + \vec{OD}.$

$$\vec{OC} = 5\vec{i} + 5\vec{j}.$$

Since D is the midpoint of AB , it has coordinates $(2, 1)$. Therefore $\vec{OD} = 2\vec{i} + \vec{j}.$

Therefore $\vec{CD} = -5\vec{i} - 5\vec{j} + 2\vec{i} + \vec{j} = -3\vec{i} - 4\vec{j}.$

b. $\vec{a} = -2\vec{i} + 4\vec{j} \quad \vec{b} = 6\vec{i} - 2\vec{j} \quad \vec{c} = 5\vec{i} + 5\vec{j}$

$$\vec{AB} = \vec{b} - \vec{a} = 8\vec{i} - 6\vec{j}$$

$$\vec{AD} = \frac{1}{2}\vec{AB} = 4\vec{i} - 3\vec{j}$$

$$\vec{CD} = -\vec{c} + \vec{a} + \vec{AD} = -3\vec{i} - 4\vec{j}$$


$$\vec{AB} \bullet \vec{CD} = (8 \times -3 + -6 \times -4) = (-24 + 24) = 0$$

Hence \vec{AB} is perpendicular to $\vec{CD}.$

QUESTION 2


$$\delta v = \pi y^2 \delta x$$

$$\begin{aligned} Vol &= \int_0^1 (\pi y^2) dx = \pi \int_0^1 (x^2 - 2x^3 + x^4) dx \\ &= \pi \left[\frac{x^3}{3} - \frac{x^4}{2} + \frac{x^5}{5} \right]_0^1 \\ &= \pi \left[\frac{1}{3} - \frac{1}{2} + \frac{1}{5} \right] - \pi[0] = \frac{\pi}{30} \text{ cubic units} \end{aligned}$$



QUESTION 3

a. $y = \log_e(2x+4)$
 $x = \log_e(2y+4)$
 $e^x = 2y+4$
 $y = \frac{e^x}{2} - 2$
 $f^{-1} : R \rightarrow R, f^{-1}(x) = \frac{e^x}{2} - 2$



$$\text{Range}(f) = R \Rightarrow \text{Dom}(f^{-1}) = R$$

b. **Y Intercept:** $f(0) = \log_e 4$
X Intercept: $\log_e(2x+4) = 0$
 $2x+4 = 1$
 $x = \frac{-3}{2}$

$$\text{Area} = \int_{\frac{-3}{2}}^0 \log_e(2x+4) dx$$

c. $y = \log_e(2x+4) \Rightarrow x = \frac{e^y}{2} - 2$

$$\begin{aligned} \int_{\frac{-3}{2}}^0 y dx &= - \int_0^{\log_e 4} x dy = \int_{\log_e 4}^0 \left(\frac{e^y}{2} - 2 \right) dy \\ &= \left[\frac{e^y}{2} - 2y \right]_{\log_e 4}^0 = \left[\frac{1}{2} \right] - \left[\frac{4}{2} - 2 \log_e 4 \right] \\ &= 2 \log_e 4 - \frac{3}{2} \text{ square units} \end{aligned}$$

QUESTION 4

a. $x = 4 \sin^3(t)$
 $y = \cos(2t)$

b. $\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx}$

$$\frac{dy}{dt} = -2 \sin(2t)$$

$$\frac{dx}{dt} = 12 \sin^2(t) \cos(t)$$

$$\frac{dt}{dx} = \frac{1}{12 \sin^2(t) \cos(t)}$$

$$\frac{dy}{dx} = \frac{-2 \sin(2t)}{12 \sin^2(t) \cos(t)} = \frac{-2 \sin(2t)}{6 \sin(t) \times 2 \sin(t) \cos(t)} = \frac{-2 \sin(2t)}{6 \sin(t) \sin(2t)}$$

$$\frac{dy}{dx} = \frac{-1}{3 \sin(t)}$$

c. $x\left(\frac{\pi}{6}\right) = 4 \sin^3\left(\frac{\pi}{6}\right) = \frac{1}{2}$

$$y\left(\frac{\pi}{6}\right) = \cos\left(2 \times \frac{\pi}{6}\right) = \frac{1}{2}$$

$$m(\text{normal}) = -\frac{dx}{dy} = 3 \sin\left(\frac{\pi}{6}\right) = \frac{3}{2}$$

Equation: Use $(y - y_1) = m(x - x_1)$

$$\left(y - \frac{1}{2}\right) = \frac{3}{2}\left(x - \frac{1}{2}\right)$$

$$y = \frac{3x}{2} - \frac{1}{4}$$

QUESTION 5

a. (i) $\frac{dh}{dt} = k, k \in R$

$$\therefore h = kt + c$$

Substitute $(0, 10)$: $c = 10$

$$\therefore h = kt + 10$$

Substitute $(5, 40)$:

$$5 = 40k + 10$$

$$k = -\frac{1}{8}$$

$$\therefore h = -\frac{t}{8} + 10$$

(ii) Substitute $t = 60$: $h = -\frac{60}{8} + 10 = -\frac{15}{2} + 10 = \frac{5}{2}$ metres.

b. (i) $\frac{dh}{dt} = -kh, k \in R^+$

$$\frac{dt}{dh} = -\frac{1}{kh}$$

$$t = -\frac{1}{k} \int \left(\frac{1}{h}\right) dh = -\frac{1}{k} \log_e h + c$$

$$\therefore h = Ae^{-kt} \text{ where } A = e^{kc}$$

Substitute $(0, 10)$: $A = 10$

$$h = 10e^{-kt}$$

Substitute $(5, 40)$: $5 = 10e^{-40k}$

$$-40k = \log_e \left(\frac{1}{2}\right)$$

$$k = \frac{1}{40} \log_e 2 = \log_e 2^{\frac{1}{40}}$$

Therefore: $h = 10e^{-t \log_e 2^{\frac{1}{40}}}$

$$h = 10e^{-kt} \text{ where } k = \frac{1}{40} \log_e 2.$$

b. (ii) From (i): $h = 10e^{-kt}$ where $k = \frac{1}{40} \log_e 2$.


Substitute $t = 60$: $h = 10e^{-60k}$ where $-60k = -\frac{60}{40} \log_e 2 = -\frac{3}{2} \log_e 2 = \log_e 2^{-3/2}$.

Therefore $h = 10e^{\log_e 2^{-3/2}} = (10)(2^{-3/2}) = \frac{10}{2^{3/2}} = \frac{10}{2\sqrt{2}} = \frac{5}{\sqrt{2}}$ metres.

QUESTION 6

a. $w = \frac{4i}{u} = \frac{4i}{-2+2i} \times \frac{-2-2i}{-2-2i} = \frac{-8i+8}{8} = 1-i$

b.



From Argand Diagram:

$$|u| = \sqrt{4+4} = 2\sqrt{2}$$

$$\text{Arg}(u) = \frac{3\pi}{4}$$

$$u = 2\sqrt{2} \text{cis}\left(\frac{3\pi}{4}\right)$$

$$|w| = \sqrt{1+1} = \sqrt{2}$$

$$\text{Arg}(w) = -\frac{\pi}{4}$$

$$w = \sqrt{2} \text{cis}\left(-\frac{\pi}{4}\right)$$

c. $4i = uw$

$$\sqrt{2} \text{cis}\left(\frac{3\pi}{4}\right) = \frac{u}{2}$$

Hence: $ua + 4i = \sqrt{2} \text{cis}\left(\frac{3\pi}{4}\right)$

$$ua + uw = \frac{u}{2}$$

$$u(a+w) = \frac{u}{2}$$

$$a+w = \frac{1}{2} \qquad a = \frac{1}{2} - w = -\frac{1}{2} + i$$

QUESTION 7

Note: $\cos \alpha = \frac{4}{5}$ and $\sin \alpha = \frac{3}{5}$.

- a. Examine particle B

$$F_N = 3mg - T = 3ma$$

$$3mg - T = \frac{3mg}{2}$$

$$T = 3mg - \frac{3mg}{2}$$

$$T = \frac{3mg}{2} \text{ Newtons}$$

- b. Examine particle A

Perpendicular to the plane:

$$N = mg \cos \alpha = \frac{4mg}{5}$$

Parallel to the plane:

$$T - \mu N - mg \sin \alpha = ma$$

$$T - \frac{4\mu mg}{5} - \frac{3mg}{5} = \frac{mg}{2}$$

$$\frac{4\mu}{5} = \frac{3}{2} - \frac{3}{5} - \frac{1}{2} = \frac{2}{5}$$

$$\mu = \frac{1}{2}$$

QUESTION 8

$$x = \cos t$$

$$\text{When } x = 0: t = \cos^{-1}(0) = \frac{\pi}{2}$$

$$\frac{dx}{dt} = -\sin t$$

$$\text{When } x = \frac{1}{2}: t = \cos^{-1}\left(\frac{1}{2}\right) = \frac{\pi}{3}$$

$$\frac{dt}{dx} = \frac{-1}{\sin t}$$

$$\frac{x^2}{\sqrt{1-x^2}} = \frac{\cos^2 t}{\sqrt{1-\cos^2 t}} = \frac{\cos^2 t}{\sqrt{\sin^2 t}} = \frac{\cos^2 t}{\sin t} = \cos^2 t \times -\frac{dt}{dx}$$

$$\text{Hence: } \int_0^{\frac{1}{2}} \frac{x^2}{(1-x^2)^{\frac{1}{2}}} dx = - \int_{t=\frac{\pi}{2}}^{t=\frac{\pi}{3}} \cos^2 t \frac{dt}{dx} dx$$

$$= \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \cos^2 t dt = \frac{1}{2} \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \cos(2t) + 1 dt$$

$$= \frac{1}{2} \left[\frac{1}{2} \sin(2t) + t \right]_{\frac{\pi}{3}}^{\frac{\pi}{2}} = \frac{1}{2} \left[\frac{1}{2} \sin \pi + \frac{\pi}{2} - \frac{1}{2} \sin \frac{2\pi}{3} - \frac{\pi}{3} \right]$$

$$= \frac{1}{2} \left[0 + \frac{3\pi}{6} - \frac{\sqrt{3}}{4} - \frac{2\pi}{6} \right]$$

$$= \frac{\pi}{12} - \frac{\sqrt{3}}{8}$$