

***INSIGHT***  
***Trial Exam Paper***

**2007**

**SPECIALIST MATHEMATICS**

**Written examination 2**

**STUDENT NAME:**

**QUESTION AND ANSWER BOOK**

**Reading time: 15 minutes**

**Writing time: 2 hours**

**Structure of book**

<i>Section</i>	<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
1	22	22	22
2	5	5	58
		Total	80

- Students are permitted to bring the following items into the examination: pens, pencils, highlighters, erasers, sharpeners, rulers, a protractor, set-squares, aids for curve sketching, once bound reference, one approved graphics calculator or CAS (memory DOES NOT need to be cleared) and, if desired, one scientific calculator.
- Students are NOT permitted to bring sheets of paper or white out liquid/tape into the examination.

**Materials provided**

- The question and answer book of 25 pages with a separate sheet of miscellaneous formulas.
- Answer sheet for multiple-choice questions

**Instructions**

- Write your **name** in the box provided and on the multiple-choice answer sheet.
- Remove the formula sheet during reading time.
- You must answer the questions in English.

**At the end of the exam**

- Place the multiple-choice answer sheet inside the front cover of this book.

**Students are NOT permitted to bring mobile phones or any other electronic devices into the examination.**

This trial examination produced by Insight Publications is NOT an official VCAA paper for the 2007 Specialist Mathematics written examination 2.

This examination paper is licensed to be printed, photocopied or placed on the school intranet and used only within the confines of the purchasing school for examining their students. No trial examination or part thereof may be issued or passed on to any other party including other schools, practising or non-practising teachers, tutors, parents, websites or publishing agencies without the written consent of Insight Publications.

Copyright © Insight Publications 2007

**SECTION 1****Instructions for Section 1**

Answer **all** questions in pencil on the multiple-choice answer sheet provided.

Choose the response that is **correct** for the question.

One mark will be awarded for a correct answer; no marks will be awarded for an incorrect answer.

Marks **are not** deducted for incorrect answers.

No marks will be awarded if more than one answer is completed for any question.

Take the **acceleration due to gravity** to have magnitude  $g \text{ m/s}^2$ , where  $g = 9.8$

**Question 1**

If  $z = i(2i + i^3 - 3)$ , then  $\text{Re}(z)$  is equal to

- A.  $-3$
- B.  $-2$
- C.  $-1$
- D.  $1$
- E.  $3$

**Question 2**

In polar form  $-\frac{1}{\sqrt{2}}(1 + i)$  is equivalent to

- A.  $\text{cis}\left(\frac{\pi}{4}\right)$
- B.  $\text{cis}\left(-\frac{3\pi}{4}\right)$
- C.  $\frac{1}{\sqrt{2}}\text{cis}\left(\frac{\pi}{4}\right)$
- D.  $\frac{1}{\sqrt{2}}\text{cis}\left(\frac{5\pi}{4}\right)$
- E.  $\frac{1}{\sqrt{2}}\text{cis}\left(-\frac{3\pi}{4}\right)$

**Question 3**

If  $u = 3 - 4i$  and  $v = 1 + 2i$  then  $\left| \frac{u^2}{v^2} \right|$  is equal to

- A.  $\bar{u}$
- B.  $-u$
- C.  $|v|$
- D.  $|v|^2$
- E.  $\left( \frac{u}{v} \right)^2$

**Question 4**


The asymptotes of the curve  $x^2 - 4y^2 + 6x + 16y = 11$  intersect at the point

- A.  $(-3, 2)$
- B.  $(-3, -8)$
- C.  $(-6, -4)$
- D.  $(3, 2)$
- E.  $(3, 8)$

**Question 5**

The range of the function  $f : \left[0, \frac{7\pi}{12}\right] \rightarrow R$ ,  $f(x) = 1 - 3\operatorname{cosec}\left(x + \frac{\pi}{4}\right)$  is

- A.  $R$
- B.  $(-\infty, -2] \cup [2, \infty)$
- C.  $(-\infty, -2]$
- D.  $[-5, -2]$
- E.  $[-5, 1 - 3\sqrt{2}]$

**Question 6**

The equation of the graph shown above could be

- A.  $y = \cos^{-1}(ax - 1)$
- B.  $y = \cos^{-1}(x - 2a)$
- C.  $y = \cos^{-1}\left(\frac{x}{2} - a\right)$
- D.  $y = \cos^{-1}\left(\frac{x-1}{a}\right)$
- E.  $y = \cos^{-1}\left(\frac{x}{a} - 1\right)$

**Question 7**

Given the function  $f(x) = \frac{|\log_e(x)|}{\log_e|x| - 1}$ , which one of the following statements is false?


- A.  $f$  has an asymptote at  $x = e$ .
- B. The point  $(1, 0) \in f(x)$ .
- C. The inverse,  $f^{-1}$ , exists for  $x \in (0, 1]$ .
- D. The range of  $f$  is  $R \setminus [0, 1)$ .
- E.  $f$  has more than two asymptotes.

**Question 8**

At the point where  $y = 2$ , the gradient of the curve  $y^2 + x^3 = 5$  is

- A.  $-3$
- B.  $-\frac{3}{4}$
- C.  $-\frac{3}{2}$
- D.  $\frac{1}{2}$
- E.  $2$

### Question 9



The graph of  $y = f(x)$  is shown above.  
Let  $F(x)$  be an antiderivative of  $f(x)$ .

The graph of  $y = F(x)$  has a


- A. local maximum at  $x = a$
- B. stationary point at  $x = b$
- C. point of inflexion at  $x = c$
- D. negative gradient for  $b < x < c$
- E. zero gradient at  $x = 0$

### Question 10

Using the substitution  $u = 2 - x$ , the integral  $\int_2^4 x(2-x)^6 dx$  is equal to

- A.  $\int_2^4 (2u^6 - u^7) du$
- B.  $\int_2^4 (u^7 - 2u^6) du$
- C.  $\int_{-2}^0 (u^7 - 2u^6) du$
- D.  $\int_0^{-2} (u^7 - 2u^6) du$
- E.  $\int_0^{-2} (2u^6 - u^7) du$

## Question 11



The shaded region shown above is formed between the graph of  $y = \sec^2(x)$  and the line joining the points  $(0, 1)$  and  $(\frac{\pi}{4}, 2)$ .

The area of the shaded region would be

- A.  $\frac{3\pi - 8}{8}$
- B.  $\frac{3\pi - 4}{8}$
- C.  $\frac{3\pi - 1}{8}$
- D.  $\frac{3\pi}{8}$
- E.  $\frac{\pi}{4}$

## Question 12

Given  $f'(x) = \frac{e^x}{e^{-x} + e^x}$  and  $f(0) = 2$ .

Using Euler's method with increments of 0.1, an approximate value of  $f(0.2)$  is

- A. 2.0500
- B. 2.0550
- C. 2.1050
- D. 2.1099
- E. 2.1484

**Question 13**

A bowl of soup is heated to a temperature of  $80^{\circ}\text{C}$ . It is then left to cool in a room in which the air temperature is  $20^{\circ}\text{C}$ . The rate at which the temperature of the soup decreases is proportional to the difference between its temperature and the temperature of the room.

Let  $S^{\circ}\text{C}$  be the temperature of the soup at any time  $t$  minutes after it is removed from the heat. Given  $k$  is a positive constant, the relationship between  $S$  and  $t$  may be modelled by the differential equation

A.  $\frac{dS}{dt} = -k(S - 20); \quad t = 0, S = 60$

B.  $\frac{dS}{dt} = -k(S - 20); \quad t = 0, S = 80$

C.  $\frac{dS}{dt} = -k(S - 60); \quad t = 0, S = 80$

D.  $\frac{dS}{dt} = -k(S - 60); \quad t = 0, S = 20$

E.  $\frac{dS}{dt} = -k(S - 80); \quad t = 0, S = 20$

**Question 14**

The solution of the differential equation  $\frac{dy}{dx} = e^{\sin(x)}$ , given  $y = 3$  when  $x = 2$ , is

A.  $y = \int_2^x e^{\sin(u)} du + 3$

B.  $y = \int_3^x e^{\sin(u)} du + 2$

C.  $y = \int_2^3 e^{\sin(x)} dx$

D.  $y = \int_2^3 e^{\sin(x)} dx + 3$

E.  $y = \int_2^3 e^{\sin(x)} dx + 2$

**Question 15**

A particle moves in a straight line such that its velocity is given by

$$v = \log_e \left| \cos\left(\frac{x}{4}\right) \right|, \quad x \in [0, 2\pi), \text{ where } x \text{ is its displacement from the origin } O.$$


The particle's acceleration  $\frac{4\pi}{3}$  units from  $O$  is

- A.  $-\frac{\sqrt{3}}{4} \log_e(2)$
- B.  $-\log_e(2)$
- C.  $-\frac{\sqrt{3}}{4}$
- D.  $\log_e(2)$
- E.  $\frac{\sqrt{3}}{4} \log_e(2)$

**Question 16**

$A$ ,  $B$  and  $C$  are three points on the circumference of a circle with centre  $O$ .

$AC$  passes through  $O$ .



Which one of the following statements is **not** true?

- A.  $\vec{AB} \cdot \vec{BC} = 0$
- B.  $\vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$
- C.  $(\vec{AB} + \vec{BC}) \cdot \vec{AC} = |\vec{AC}|^2$
- D.  $\vec{AB} \cdot \vec{AC} = |\vec{AB}| |\vec{AC}| \cos(A)$
- E.  $|\vec{AC}| = |\vec{AB}| + |\vec{BC}|$

**Question 17**

Let  $\underline{m} = 4\underline{i} - \underline{j} + 2\underline{k}$  and  $\underline{n} = \underline{i} + \underline{j} - 2\underline{k}$ .

A unit vector in the direction of  $\underline{m} - 2\underline{n}$  would be

- A.  $\frac{1}{7}(2\underline{i} - 3\underline{j} + 6\underline{k})$
- B.  $\frac{1}{3}(2\underline{i} + \underline{j} + 2\underline{k})$
- C.  $\frac{1}{\sqrt{17}}(2\underline{i} - 3\underline{j} - 2\underline{k})$
- D.  $\frac{1}{\sqrt{41}}(6\underline{i} + \underline{j} + 4\underline{k})$
- E.  $\frac{1}{\sqrt{29}}(3\underline{i} - 2\underline{j} + 4\underline{k})$

**Question 18**


The velocity of a particle at time  $t$ ,  $t \geq 0$  is given by  $\underline{v} = 4 \sin(2t)\underline{i} + 6 \cos(3t)\underline{j}$ .

If the particle was initially at  $\underline{i} + 2\underline{j}$ , its position after  $\frac{\pi}{2}$  seconds will be

- A.  $-3\underline{i} + 4\underline{j}$
- B.  $3\underline{i} - 2\underline{j}$
- C.  $3\underline{i} + 3\underline{j}$
- D.  $5\underline{i}$
- E.  $5\underline{i} - \underline{j}$

**Question 19**

The graph below shows the velocity,  $v$  m/s, of a particle moving in a straight line for 25 s.




How many metres is the particle from its starting point after 25 s?

- A. 6
- B. 14
- C. 30
- D. 46
- E. 50

**Question 20**

Two forces act simultaneously on a particle, as shown in the diagram below. One force of 9 N acts due west and another force of 11 N acts at an angle of  $60^\circ$  in an anticlockwise direction from due east.




Correct to the nearest degree, the resultant force acting in an anticlockwise direction from due east will be

- A.  $50^\circ$
- B.  $70^\circ$
- C.  $110^\circ$
- D.  $120^\circ$
- E.  $130^\circ$


**Question 21**

A mass of 25 kg is pulled across a smooth horizontal surface by a force of 30 newtons acting at an angle of  $20^\circ$  to the horizontal level.



The magnitude of the normal reaction of the surface on the mass, in newtons, is closest to

- A. 215
- B. 217
- C. 235
- D. 245
- E. 255

**Question 22**

Two bodies each of mass,  $m$  kg, are connected on a back-to-back plane by a light string passing over a smooth pulley, as shown in the diagram. The coefficient of friction of the surface of each plane is  $\mu$ .

If the system is on the point of moving in the direction shown, the value of  $\mu$  will be

- A.  $\tan(\theta)$
- B.  $\cot(\theta)$
- C.  $\sin(\theta) - \cos(\theta)$
- D.  $\frac{\sin(\theta) - \cos(\theta)}{\sin(\theta) + \cos(\theta)}$
- E.  $\frac{\cos(\theta) - \sin(\theta)}{\cos(\theta) + \sin(\theta)}$

**END OF SECTION A  
TURN OVER**

## SECTION 2

## Instructions for Section 2

Answer all the questions in the spaces provided.

A decimal approximation will not be accepted if an exact answer is required to a question.

In questions where more than one mark is available, approximate working must be shown.


Unless otherwise indicated, the diagrams in this book have not been drawn to scale.

Take the **acceleration due to gravity** to have magnitude  $g \text{ m/s}^2$ , where  $g = 9.8$

## Question 1

Let  $u = 2\text{cis}\left(\frac{\pi}{3}\right)$ .

- a. i.  $u$ ,  $\bar{u}$  and  $v$  are solutions of the equation  $\{z : z^3 = k, z \in C\}$ .  
Plot  $u$ ,  $\bar{u}$  and  $v$  on the Argand plane below.



3 marks

- ii. Determine the value of  $k$ .

---

1 mark

**b.** Show that  $u \in \{z : \operatorname{Re}(z) + \sqrt{3} \operatorname{Im}(z) = 4\}$ .

---



---



---



---



---



---

2 marks

**c. i.** Sketch  $\{z : \operatorname{Re}(z) + \sqrt{3} \operatorname{Im}(z) = 4\}$  on the Argand plane given in part **a**, showing the exact intercepts with the axes.

1 mark

**ii. Hence,** shade the region represented by

$$\left\{z : \frac{\pi}{6} \leq \operatorname{Arg}(z) \leq \frac{\pi}{3}\right\} \cap \{z : \operatorname{Re}(z) + \sqrt{3} \operatorname{Im}(z) \leq 4\}.$$

1 mark

**iii.** Calculate the exact area of this region.

---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---

3 marks

Total 3 + 1 + 2 + 1 + 1 + 3 = 11 marks

**SECTION B** – continued  
**TURN OVER**

**Question 2**

Given  $f : R \rightarrow R$ ,  $f(x) = \frac{x}{x^2 + 2} + 1$

**a. i.** Show that  $f'(x) = \frac{2 - x^2}{(x^2 + 2)^2}$ .

---



---



---

1 mark

**ii. Hence,** determine the exact coordinates of any stationary points of  $f$ .

---



---



---



---



---

2 marks

**b. i.** Use calculus to show that  $f$  has points of inflexion.

---



---



---



---



---



---



---



---



---




---

2 marks



**Question 3**

The tank shown below has the shape of a truncated cone with base diameter 1.4 m, top diameter 1 m and height 2 m. It may be modelled by rotating the line segment  $MN$  around the  $y$ -axis.



- a. The line segment  $MN$  has equation  $ax + by + c = 0$ . Show that  $a = 10$ ,  $b = 1$  and  $c = -7$ .

---



---



---



---



---



---



---



---


2 marks





**Question 4**

A yacht, initially at point  $B$ , will sail to point  $P(2, 7)$  on the other side of the bay. Distances are measured in kilometres in relation to the origin,  $O$ .



- a. Write a vector  $\vec{OP}$  that gives the position of point  $P$ .

---



---

1 mark

The yacht leaves  $B$ , sailing with a velocity of  $\underline{v} = -2\underline{i} + 2t\underline{j}$  km/h. It reaches  $P$  after 3 h.

- b. i. Determine the speed of the yacht when it reaches  $P$ .  
Write your answer correct to 1 decimal place.

---



---



---

1 mark


**SECTION B – continued**  
**TURN OVER**





**Question 5**

A wire rope passing over a smooth pulley is used to transport materials to and from a building site situated below ground level. A 1 tonne engine at ground level applies a horizontal force of  $P$  newton along a track to pull a load of  $L$  kg upwards. The coefficient friction between the engine and the track is 0.25.



- a.** The load,  $L$  kg, is on the point of moving.  
**i.** On the diagram above, show all forces acting.

1 mark

- ii.** Find  $P$  in terms of  $L$ .

---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---

4 marks



- ii. Determine the time taken, in seconds, to raise the load from rest to a point 15 m above the level of the building site.

---



---



---




---

1 mark

When the engine reaches the end of the track it applies its breaks so that the load will remain stationary at the loading platform.

The diagram below shows a 300 kg load suspended at the loading platform 40 m above the building site. It is stationary.



- c. Find the minimum force that the engine's breaks need to apply in order to keep the 300 kg load stationary in this position.

---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---

3 marks

**SECTION B** – continued

