

Year 2006
VCE
Specialist Mathematics
Solutions
Trial Examination 2



KILBAHA MULTIMEDIA PUBLISHING
PO BOX 2227
KEW VIC 3101
AUSTRALIA

TEL: (03) 9817 5374
FAX: (03) 9817 4334
chemas@chemas.com
www.chemas.com

© Kilbaha Pty Ltd 2006

ABN 47 065 111 373

© KILBAHA PTY LTD 2006

IMPORTANT COPYRIGHT NOTICE

- This material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Pty Ltd.
 - The contents of this work are copyrighted. Unauthorised copying of any part of this work is illegal and detrimental to the interests of the author.
 - For authorised copying within Australia please check that your institution has a licence from Copyright Agency Limited. This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.
 - Teachers and students are reminded that for the purposes of school requirements and external assessments, students must submit work that is clearly their own.
 - Schools which purchase a licence to use this material may distribute this electronic file to the students at the school for their exclusive use. This distribution can be done either on an Intranet Server or on media for the use on stand-alone computers.
 - Schools which purchase a licence to use this material may distribute this printed file to the students at the school for their exclusive use.
-
- **The Word file (if supplied) is for use ONLY within the school.**
 - **It may be modified to suit the school syllabus and for teaching purposes.**
 - **All modified versions of the file must carry this copyright notice.**
 - **Commercial use of this material is expressly prohibited.**

SECTION 1

ANSWERS

1	A	B	C	D	E
2	A	B	C	D	E
3	A	B	C	D	E
4	A	B	C	D	E
5	A	B	C	D	E
6	A	B	C	D	E
7	A	B	C	D	E
8	A	B	C	D	E
9	A	B	C	D	E
10	A	B	C	D	E
11	A	B	C	D	E
12	A	B	C	D	E
13	A	B	C	D	E
14	A	B	C	D	E
15	A	B	C	D	E
16	A	B	C	D	E
17	A	B	C	D	E
18	A	B	C	D	E
19	A	B	C	D	E
20	A	B	C	D	E
21	A	B	C	D	E
22	A	B	C	D	E

SECTION 1

Question 1 **Answer D**

The graphs of all alternatives give vertical asymptotes at $x=0$, $\frac{\pi}{2}$ and π however option D is wrong, it is the reflection in the x -axis

Question 2 **Answer D**

$$f(g(x)) = \frac{1}{-x^2 - (a-b)x + ab} = \frac{1}{-(x^2 + (a-b)x - ab)} = \frac{1}{-(x-b)(x+a)}$$

So there are vertical asymptotes at $x=-a$ and $x=b$

Question 3 **Answer A**

$$y = \frac{ax^2 + b}{x} = ax + \frac{b}{x} = ax + bx^{-1}$$

$$\frac{dy}{dx} = a - bx^{-2} = a - \frac{b}{x^2} \quad \text{for turning points } \frac{dy}{dx} = 0$$


$$a = \frac{b}{x^2} \quad x^2 = \frac{b}{a} \quad x = \pm \sqrt{\frac{b}{a}} \quad \text{however there are no turning points, so there is no}$$

solutions for $x = \pm \sqrt{\frac{b}{a}}$ so a and b must have opposite signs the product $ab < 0$ so $a > 0$ and $b < 0$

Question 4 **Answer B**

$$\operatorname{cosec}(x) = \frac{1}{\sin(x)} = \frac{4\sqrt{7}}{7}$$

$$\sin(x) = \frac{7}{4\sqrt{7}} = \frac{\sqrt{7}}{4}$$



so from the triangles $\cos(x) = -\frac{3}{4}$ since $\frac{\pi}{2} < x < \pi$ x is obtuse and in the second

$$\text{quadrant } \sec(x) = \frac{1}{\cos(x)} = -\frac{4}{3}$$

Question 5**Answer E**

The ellipse $\frac{(y+1)^2}{18} + \frac{(x-4)^2}{8} = 2$ in standard form is $\frac{(x-4)^2}{16} + \frac{(y+1)^2}{36} = 1$
 it has centre at $(4, -1)$ $a = 4$ $b = 6$ the domain is $4 \pm 4 = [0, 8]$
 and the range $-1 \pm 6 = [-7, 5]$

Question 6**Answer C**

Given $u = 3 \operatorname{cis}\left(\frac{\pi}{4}\right)$, $v = a \operatorname{cis}(b)$ and $\frac{u}{v} = \frac{3 \operatorname{cis}\left(\frac{\pi}{4}\right)}{a \operatorname{cis}(b)} = \frac{3}{a} \operatorname{cis}\left(\frac{\pi}{4} - b\right) = -6 = 6 \operatorname{cis}(\pi)$
 so $\frac{3}{a} = 6$ $a = \frac{1}{2}$ and $\frac{\pi}{4} - b = \pi$ $b = -\frac{3\pi}{4}$

Question 7**Answer E**

$z = (\cos \theta + i \sin \theta)^3 = \operatorname{cis}(3\theta)$ by DeMoivre's Theorem
 $z^2 = (\cos \theta + i \sin \theta)^6 = \operatorname{cis}(6\theta) = \cos(6\theta) + i \sin(6\theta)$ and $z^2 = a + bi$
 Equating real and imaginary parts respectively gives $\cos(6\theta) = a$ and $\sin(6\theta) = b$

Question 8**Answer C**

The circle has radius 2, one of the solutions is $u = 2i$ and $u^3 = 8i^3 = -8i$
 but $P(u) = 0$ so $P(z) = z^3 + 8i = 0$

Question 9**Answer A**

$P(z) = z^3 + bz^2 + cz + d$ and $P(-\alpha i) = 0$ and $P(\beta) = 0$ by the conjugate root theorem $P(\alpha i) = 0$ so $z^2 + \alpha^2$ is one factor, the other is $(z - \beta)$ expanding
 $P(z) = (z^2 + \alpha^2)(z - \beta) = z^3 - \beta z^2 + \alpha^2 z - \alpha^2 \beta = z^3 + bz^2 + cz + d = 0$ equating
 $b = -\beta$ $c = \alpha^2$ $d = -\alpha^2 \beta$

Question 10**Answer C**

The inside of the circle of radius 4, can be represented by $\{z: |z| \leq 4\}$ or $\{z: z\bar{z} \leq 16\}$ or $\{z: \operatorname{Re}^2(z) + \operatorname{Im}^2(z) \leq 16\}$ the upper half plane of the Argand diagram can be represented by $\{z: \operatorname{Im}(z) \geq 0\}$ or $\{z: 0 \leq \operatorname{Arg}(z) \leq \pi\}$ option C is the only correct intersection of all the possible correct representations.

Question 11**Answer E**

When we have the repeated factor $(x+3)^2$ and the non-linear factor (x^2+3) we represent the expression $\frac{3x+1}{(x+3)^2(x^2+3)}$ in partial fractions as

$$\frac{A}{x+3} + \frac{B}{(x+3)^2} + \frac{Cx+D}{x^2+3}$$

Question 12**Answer E**

$$\int_1^4 \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = \int_1^4 \cos(\sqrt{x}) \frac{1}{\sqrt{x}} dx$$

$$\text{let } u = \sqrt{x} = x^{\frac{1}{2}} \quad \frac{du}{dx} = \frac{1}{2} x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}} \quad \frac{1}{\sqrt{x}} dx = 2 du$$

change the terminals when $x=4$ $u=2$ and when $x=1$ $u=1$


the definite integral becomes $2 \int_1^2 \cos(u) du$

Question 13**Answer D**

At $x=a$ and $x=e$ the gradient changes from negative to positive, this gives a minimum at the two points $x=a$ and $x=e$. At $x=c$ the gradient changes from positive to negative this gives a maximum at the point $x=c$. The graph of $F(x)$ is a quartic and will have two points of inflexions, one somewhere between $x=a$ and $x=c$ and another somewhere between $x=c$ and $x=e$.

Question 14

Answer C



Option A is true $\overrightarrow{AP} = \overrightarrow{PB} = \frac{1}{2} \overrightarrow{AB}$ since P is the mid-point of \overline{AB}

Option B is true since ABC is an equilateral triangle all side lengths are equal

$$|\overline{AB}| = |\overline{BC}| = |\overline{AC}|$$

Option C is false \overrightarrow{AP} is perpendicular to \overline{PC} so $\overrightarrow{AP} \cdot \overline{PC} = 0$

Option D is true the angle CAB is 60° $\cos(60^\circ) = \frac{1}{2} = \frac{\overline{AC} \cdot \overline{AB}}{|\overline{AC}| |\overline{AB}|}$

Option E is true the angle PCB is 30° $\cos(30^\circ) = \frac{\sqrt{3}}{2} = \frac{\overline{CP} \cdot \overline{CB}}{|\overline{CP}| |\overline{CB}|}$

Question 15

Answer A

$$\underline{a} = -2\underline{i} + 2\underline{j} - \underline{k} \quad \text{and} \quad \underline{b} = 4\underline{i} + y\underline{j} + 2\underline{k} \quad \underline{a} \cdot \underline{b} = -8 + 2y - 2 = 2y - 10$$

$$|\underline{a}| = \sqrt{4 + 4 + 1} = 3 \quad |\underline{b}| = \sqrt{16 + y^2 + 4} = \sqrt{20 + y^2}$$

So the angle between \underline{a} and \underline{b} is $\cos^{-1} \left(\frac{2y - 10}{3\sqrt{(20 + y^2)}} \right)$

If $y = -4$ then $-2\underline{a} = \underline{b}$ so the vectors \underline{a} and \underline{b} are linearly **dependent**, option A is false

All the other alternatives are true.


Option C if \underline{a} is perpendicular to \underline{b} then $2y - 10 = 0 \quad y = 5$

Option D if $y > 5$ then the angle between the vectors \underline{a} and \underline{b} is acute.

Option E if $y < 5$ then the angle between the vectors \underline{a} and \underline{b} is obtuse.

Question 16**Answer B**

The velocity vector is $\dot{r}(t) = 4e^{\frac{t}{2}}\underline{i} - 2\sin\left(\frac{t}{2}\right)\underline{j}$ integrating with respect to t to get the position vector $r(t) = \int 4e^{\frac{t}{2}} dt \underline{i} - \int 2\sin\left(\frac{t}{2}\right) dt \underline{j} = 8e^{\frac{t}{2}}\underline{i} + 4\cos\left(\frac{t}{2}\right)\underline{j} + \underline{C}$ now using the initial conditions $r(0) = 3\underline{i} - 3\underline{j} = 8\underline{i} + 4\underline{j} + \underline{C}$ so $\underline{C} = -5\underline{i} - 7\underline{j}$ and

$$r(t) = \left(8e^{\frac{t}{2}} - 5\right)\underline{i} + \left(4\cos\left(\frac{t}{2}\right) - 7\right)\underline{j}$$
Question 17**Answer A**

Option A is false if $\theta = 45$ then $P = Q = \frac{\sqrt{2}R}{2}$


All the other alternatives are true

Option B $R^2 = P^2 + Q^2$

Option C $P = R \sin(\theta)$ and $Q = R \cos(\theta)$

Option D $\tan(\theta) = \frac{P}{Q}$

Option E as vectors $\underline{P} + \underline{Q} + \underline{R} = \underline{0}$

Question 18**Answer E**


$$m = 60\text{kg} \quad g = 9.8\text{m/s}^2 \quad \mu = 0.75$$

$$N = mg = 588 \text{ newtons}$$

$$\mu N = 441 \text{ newtons}$$

movement with constant acceleration

is only possible if $F > \mu N$

Question 19**Answer B**

resolving the forces, perpendicular to the plane

$$N - mg \cos(\theta) = 0 \Rightarrow N = mg \cos(\theta)$$

resolving the forces up and parallel to the plane.

$$P + \mu N - mg \sin(\theta) = 0$$

$$P + \mu mg \cos(\theta) - mg \sin(\theta) = 0$$

$$P = mg \sin(\theta) - \mu mg \cos(\theta)$$

$$P = mg (\sin \theta - \mu \cos \theta)$$

Question 20**Answer A**

$$\tan^{-1}\left(\frac{1}{2t}\right) \frac{dt}{dy} = 1 \quad \frac{dy}{dt} = \tan^{-1}\left(\frac{1}{2t}\right) \text{ using a calculator program gives}$$

t	y
1.00	2.0000
1.25	2.1159
1.50	2.2110

Question 21**Answer B**

The solution curves are of the form $x = \sin(2t) + c$ so differentiating,

the differential equation is $\frac{dx}{dt} = 2 \cos(2t)$

Question 22**Answer B**

The volume in the tank is $V = A h$ $\frac{dV}{dh} = A$

and $\frac{dV}{dt} = \text{inflow} - \text{outflow} = Q - c\sqrt{h}$


$\frac{dh}{dt} = \frac{dh}{dV} \frac{dV}{dt}$ so that $\frac{dh}{dt} = \frac{Q - c\sqrt{h}}{A}$ $h(0) = h_0$

END OF SECTION 1 SUGGESTED ANSWERS

SECTION 2

Question 1

- a.i.** $3x^2 + 18x - y^2 + 4y + 11 = 0$
 $3(x^2 + 6x) - (y^2 - 4y) = -11$
 $3(x^2 + 6x + 9) - (y^2 - 4y + 4) = -11 + 27 - 4$ M1
 $3(x+3)^2 - (y-2)^2 = 12$ $\frac{(x+3)^2}{4} - \frac{(y-2)^2}{12} = 1$ A1
- ii.** centre (h, k) where $h = -3$ $k = 2$ $a = 2$ $b = 2\sqrt{3}$
 hyperbola, centre $(-3, 2)$ vertices $(-5, 2)$ $(-1, 2)$ A1
 asymptotes $\frac{y-2}{2\sqrt{3}} = \pm \frac{x+3}{2}$ $y = \pm\sqrt{3}(x+3) + 2$
 $y = \sqrt{3}x + 2 + 3\sqrt{3}$ and $y = -\sqrt{3}x + 2 - 3\sqrt{3}$ A1



- b.i** $3x^2 + 18x - y^2 + 4y + 11 = 0$
 differentiating implicitly
 $\frac{d}{dx}(3x^2) + \frac{d}{dx}(18x) - \frac{d}{dx}(y^2) + \frac{d}{dx}(4y) + \frac{d}{dx}(11) = 0$
 $6x + 18 - 2y \frac{dy}{dx} + 4 \frac{dy}{dx} = 0$
 $\frac{dy}{dx}(2y - 4) = 6x + 18$
 $\frac{dy}{dx} = \frac{3x + 9}{y - 2} = \frac{3(x + 3)}{y - 2}$ A1

- ii. When the tangent is parallel to the y-axis. $\frac{dy}{dx} \rightarrow \infty \Rightarrow y-2=0 \quad y=2$

These are obviously the vertices $(-5, 2)$ $(-1, 2)$ A1

- c. $R(0, 2-3\sqrt{3})$, $S(0, 3\sqrt{3}+2)$ and $C(-3, 2)$

$$\overline{CS} = \overline{OS} - \overline{OC} = 3\hat{i} + 3\sqrt{3}\hat{j} \quad \overline{CR} = \overline{OR} - \overline{OC} = 3\hat{i} - 3\sqrt{3}\hat{j} \quad \text{M1}$$

$$|\overline{CS}| = \sqrt{9+27} = 6 \quad |\overline{CR}| = \sqrt{9+27} = 6$$

$$\overline{CS} \cdot \overline{CR} = 9 - 9 \times 3 = -18$$

The angle between \overline{CS} and \overline{CR} is θ where $\cos(\theta) = \frac{\overline{CS} \cdot \overline{CR}}{|\overline{CS}| |\overline{CR}|} = \frac{-18}{36} = -\frac{1}{2}$

$$\theta = \cos^{-1}\left(-\frac{1}{2}\right) = 120^\circ \quad \text{A1}$$

this angle represents the angle between the asymptotes at the centre of the hyperbola. A1

- d. $m = -4 + yi$, $f = -7 + 2i$ and $z = x + yi$

$$|z - f| = 2 |z - m|$$

$$|(x + yi) - (-7 + 2i)| = 2 |(x + yi) - (-4 + yi)|$$

$$|(x + 7) + (y - 2)i| = 2 |(x + 4) + (y - y)i| \quad \text{M1}$$

$$\sqrt{(x + 7)^2 + (y - 2)^2} = 2\sqrt{(x + 4)^2} \quad \text{squaring both sides and expanding} \quad \text{M1}$$

$$x^2 + 14x + 49 + y^2 - 4y + 4 = 4(x^2 + 8x + 16) = 4x^2 + 32x + 64$$

$$3x^2 + 18x - y^2 + 4y + 11 = 0$$

$$\frac{(x + 3)^2}{4} - \frac{(y - 2)^2}{12} = 1 \quad \text{the hyperbola again} \quad \text{A1}$$

- e. $r(t) = (-3 + 2\sec(2t))\hat{i} + (2 + 2\sqrt{3}\tan(2t))\hat{j}$

$$x = -3 + 2\sec(2t) \quad y = 2 + 2\sqrt{3}\tan(2t)$$

$$\sec(2t) = \frac{x + 3}{2} \quad \tan(2t) = \frac{y - 2}{2\sqrt{3}} \quad \text{M1}$$

$$\sec^2(2t) - \tan^2(2t) = 1$$


$$\frac{(x + 3)^2}{4} - \frac{(y - 2)^2}{12} = 1 \quad \text{the hyperbola again} \quad \text{A1}$$

Question 2

a. domain $|x| \leq 15 = [-15, 15]$

A1

b.



A1


c. $f(x) = 25 - 20 \cos^{-1}\left(\frac{x}{15}\right) \quad x \in [-15, 15]$

end points $(-15, 25 - 20\pi)$ $(15, 25)$

$$f'(x) = \frac{20}{\sqrt{225 - x^2}} \quad x \in (-15, 15)$$

the function is defined at the end-points
but the gradient is not defined at the end-points.

A1



- d.** when $y = 0$ $25 - 20 \cos^{-1}\left(\frac{x}{15}\right) = 0 \Rightarrow x = 4.72984$
 the diameter of the base of the vase is $2 \times 4.72984 = 9.460$ cm A1
- e.** $f'(10) = \frac{20}{\sqrt{225-100}} = \frac{4\sqrt{5}}{5} \approx 1.78885 = \tan(\theta)$ M1
 $\theta = \tan^{-1}(1.789) = 60.794^\circ = 60^\circ 48'$ A1
- f. i.** $y = 25 - 20 \cos^{-1}\left(\frac{x}{15}\right)$
 $20 \cos^{-1}\left(\frac{x}{15}\right) = 25 - y$
 $\cos^{-1}\left(\frac{x}{15}\right) = \frac{25 - y}{20}$ M1
 $\frac{x}{15} = \cos\left(\frac{25 - y}{20}\right)$
 $x = 15 \cos\left(\frac{25 - y}{20}\right)$ M1
 $V_y = \pi \int_a^b x^2 dy$
 $V = 225\pi \int_0^{25} \cos^2\left(\frac{25 - y}{20}\right) dy$ A1
- ii.** $V = 225\pi \int_0^{25} \cos^2\left(\frac{25 - y}{20}\right) dy = 10950.90 \text{ cm}^3$ A1

Question 3

a. $45a = -260\sqrt{v}$
 $9a = -52\sqrt{v}$ A1

b. $a = \frac{dv}{dt} = -\frac{52\sqrt{v}}{9}$ inverting both sides
 $\frac{dt}{dv} = -\frac{9}{52\sqrt{v}}$
 $\int \frac{dv}{\sqrt{v}} = -\frac{52}{9} \int 1 dt$ M1

$$\int v^{-\frac{1}{2}} dv = -\frac{52t}{9} + C_1$$

$$2\sqrt{v} = -\frac{52t}{9} + C_1 \quad \text{now when } t=0 \quad v=9$$

$$2\sqrt{9} = C_1 = 6$$

$$2\sqrt{v} = 6 - \frac{52t}{9} \quad \text{now when } v=1$$

$$2 = 6 - \frac{52t}{9}$$


$$\frac{52t}{9} = 4$$

$$t = \frac{36}{52} = \frac{9}{13}$$

The head-wind blows for $\frac{9}{13}$ secs A1

c.i. $\sqrt{v} = 3 - \frac{26t}{9}$
 $v = v(t) = \left(3 - \frac{26t}{9}\right)^2 = \frac{(27 - 26t)^2}{81}$ A1

ii.



Note the restricted domain for t as $\left[0, \frac{9}{13}\right]$, graph passes through the points

$$(0, 9) \text{ and } \left(\frac{9}{13}, 1\right) \quad \text{A1}$$

d. Since the area under the velocity time graph represents the displacement

$$D = \int_0^{\frac{9}{13}} \left(3 - \frac{26t}{9}\right)^2 dt = 3 \quad \text{A1}$$

Pete rode his bike exactly 3 metres during the head-wind. A1

e. $45a = -260\sqrt{v}$ use $a = v \frac{dv}{dx}$

$$45v \frac{dv}{dx} = -260\sqrt{v}$$

$$\frac{dv}{dx} = -\frac{52}{9\sqrt{v}} \quad \text{A1}$$

f. $\frac{dv}{dx} = -\frac{52}{9\sqrt{v}}$ inverting both sides


$$\frac{dx}{dv} = -\frac{9\sqrt{v}}{52} \quad \text{integrating with respect to } v \quad \text{M1}$$

$$x = -\frac{9}{52} \int_9^1 \sqrt{v} dv = \frac{9}{52} \int_1^9 \sqrt{v} dv \quad \text{A1}$$

$$D = \frac{9}{52} \left[\frac{2}{3} v^{\frac{3}{2}} \right]_1^9 = \frac{3}{26} (27 - 1) = 3 \quad \text{A1}$$

Question 4

a. i.



ii Using Newtons 2nd Law of Motion

resolving horizontally around the log in the \hat{i} direction.

$$(1) \quad T \cos(35^\circ) - 0.8N = 400 \times 0.1$$

resolving vertically around the log in the \hat{j} direction.

$$(2) \quad T \sin(35^\circ) + N - 400g = 0$$

$$(2) \quad \Rightarrow N = 400g - T \sin(35^\circ) \text{ into (1)}$$

$$T \cos(35^\circ) - 0.8(400g - T \sin(35^\circ)) = 40$$

$$T(\cos(35^\circ) + 0.8 \sin(35^\circ)) - 0.8 \times 400g = 40$$


$$T = \frac{40(1 + 10 \times 0.8 \times 9.8)}{\cos(35^\circ) + 0.8 \sin(35^\circ)} = 2485.11 = 2485 \text{ newtons}$$

iii. resolving horizontally around the elephant in the \hat{i} direction.

$$(3) \quad P - T \cos(35^\circ) = 5000 \times 0.1$$

$$P = T \cos(35^\circ) + 500 = 2535.68$$

$$P = 2536 \text{ newtons}$$



- ii.** constant speed is zero acceleration. Note the 0.1 m/s is not used.
resolving up and parallel to the hill around the log in the \hat{i} direction.

$$(1) T \cos(35^\circ) - 0.8N - 400g \sin(5^\circ) = 0$$

resolving perpendicular to the hill around the log in the \hat{j} direction

$$(2) N + T \sin(35^\circ) - 400g \cos(5^\circ) = 0 \quad \text{M1}$$

$$(2) \Rightarrow N = 400g \cos(5^\circ) - T \sin(35^\circ) \quad \text{into (1)}$$

$$T \cos(35^\circ) - 0.8(400g \cos(5^\circ) - T \sin(35^\circ)) - 400g \sin(5^\circ) = 0$$

$$T(\cos(35^\circ) + 0.8 \sin(35^\circ)) = 400g(\sin(5^\circ) + 0.8 \cos(5^\circ)) \quad \text{M1}$$

$$T = \frac{400g(\sin(5^\circ) + 0.8 \cos(5^\circ))}{\cos(35^\circ) + 0.8 \sin(35^\circ)} = 2711.801$$

$$T = 2712 \text{ newtons} \quad \text{A1}$$

- iii.** resolving horizontally around the elephant in the \hat{i} direction.

$$(3) Q - T \cos(35^\circ) - 5000g \sin(5^\circ) = 0$$

$$Q = T \cos(35^\circ) + 5000g \sin(5^\circ) = 6492.01$$

$$Q = 6492 \text{ newtons} \quad \text{A1}$$

Question 5 $\underline{r}(t) = (-25t^2 + 52.5t) \underline{i} + \left(2e^{-\frac{7t}{2}} \left| \cos\left(\frac{17\pi t}{10}\right) \right| \right) \underline{j} \quad t \geq 0$

a. $\underline{r}(0) = 0 \underline{i} + 2 \underline{j} = 2 \underline{j}$

The ball was 2 metres above the ground when it left the bowlers hand. A1

b. differentiating using the product in the \underline{j} component M1

$$\dot{\underline{r}}(t) = (-50t + 52.5) \underline{i} + \left(-\frac{7}{2} \times 2e^{-\frac{7t}{2}} \left| \cos\left(\frac{17\pi t}{10}\right) \right| + 2e^{-\frac{7t}{2}} \times -\frac{17\pi}{10} \left| \sin\left(\frac{17\pi t}{10}\right) \right| \right) \underline{j}$$

$$\dot{\underline{r}}(t) = (-50t + 52.5) \underline{i} - \left(e^{-\frac{7t}{2}} \left(7 \left| \cos\left(\frac{17\pi t}{10}\right) \right| + \frac{17\pi}{5} \left| \sin\left(\frac{17\pi t}{10}\right) \right| \right) \right) \underline{j} \quad \text{A1}$$

c. $\dot{\underline{r}}(0) = 52.5 \underline{i} - 7 \underline{j} \quad |\dot{\underline{r}}(0)| = \sqrt{52.5^2 + (-7)^2} = 53.0 \text{ m/s} \quad \text{M1}$

$$m = 155 \text{ gm} = 0.155 \text{ kg}$$

magnitude of the momentum of the ball when it left the bowlers hand

$$|\underline{p}| = m |\dot{\underline{r}}(0)| = 0.155 \times 53.0 = 8.21 \text{ kg m/s} \quad \text{A1}$$

d. The ball hits the ground when $2e^{-\frac{7t}{2}} \left| \cos\left(\frac{17\pi t}{10}\right) \right| = 0$

$$\cos\left(\frac{17\pi t}{10}\right) = 0 \quad \text{M1}$$

$$\frac{17\pi t}{10} = \frac{\pi}{2}$$

$$t = \frac{5}{17} \text{ sec} \quad \text{hits the ground after } 0.294 \text{ seconds} \quad \text{A1}$$

$$\underline{r}\left(\frac{5}{17}\right) = \left(-25\left(\frac{5}{17}\right)^2 + 52.5 \times \frac{5}{17} \right) \underline{i} + \left(2e^{-\frac{7 \times \frac{5}{17}}{2}} \left| \cos\left(\frac{17\pi}{10} \times \frac{5}{17}\right) \right| \right) \underline{j}$$


$$\underline{r}\left(\frac{5}{17}\right) = 13.278 \underline{i} + 0 \underline{j}$$

$$\underline{r}\left(\frac{5}{17}\right) = 13.278 \underline{i} \quad (20 - 13.278 = 6.72)$$

The ball strikes the ground 6.72 metres from the batting stumps A1

e.
$$\dot{\underline{r}}\left(\frac{5}{17}\right) = \left(-50 \times \frac{5}{17} + 52.5\right) \underline{i} + \left(e^{-\frac{7.5}{17}} \left(-7 \left|\cos\left(\frac{\pi}{2}\right)\right| - \frac{17\pi}{5} \left|\sin\left(\frac{\pi}{2}\right)\right|\right)\right) \underline{j}$$

$$\dot{\underline{r}}\left(\frac{5}{17}\right) = 37.794 \underline{i} - 3.8156 \underline{j}$$



M1

$$\tan(\theta) = \frac{\dot{y}}{\dot{x}} = \frac{3.8156}{37.794}$$

$$\theta = \tan^{-1}(0.101) = 5.765^\circ = 5^\circ 46'$$

The ball strikes the ground at an angle of $5^\circ 46'$

A1

f. when $x = 20$ $20 = -25t^2 + 52t$

$$25t^2 - 52.5t + 20 = 0$$

$$t = \frac{1}{2} = 0.5 \text{ seconds}$$

Hits the batting stumps after 0.5 seconds

A1

$$\underline{r}\left(\frac{1}{2}\right) = 20 \underline{i} + \left(2e^{-\frac{7 \times 0.5}{2}} \left|\cos\left(\frac{17\pi}{20}\right)\right|\right) \underline{j}$$

$$\underline{r}\left(\frac{1}{2}\right) = 20 \underline{i} + 0.3097 \underline{j}$$

hits the stumps 31 cm above the ground.

A1

g. Total distance of 20 metres in 0.5 seconds, gives the average

velocity of the ball as 40 m/s or $40 \times \frac{60 \times 60}{1000} = 144 \text{ km/hr}$

A1

END OF SECTION 2 SUGGESTED ANSWERS